arXiv:1701.06475v1 [math.AC] 23 Jan 2017

MODULES WITH PURE RESOLUTIONS

H. ANANTHNARAYAN AND RAJIV KUMAR

Abstract. Let R a standard graded algebra over a field k. In this paper, we show that the property of R being Cohen-Macaulay is characterized by the existence of a pure Cohen-Macaulay R-module corresponding to any degree sequence of length at most $\operatorname{depth}(R)$. We also give a relation in terms of graded Betti numbers, called the Herzog-Kühl equations, for a pure R-module M to satisfy the condition $\dim(R) - \operatorname{depth}(R) = \dim(M) - \operatorname{depth}(M)$. When R is Cohen-Macaulay, we prove an analogous result characterizing all graded Cohen-Macaulay R-modules.

Introduction

Let $S = \mathsf{k}[X_1, \ldots, X_n]$ be a polynomial ring over a field k. M. Boij and J. Söderberg conjectured([1, Conjecture 2.4]) that given any degree sequence $\mathbf{d}$ of length at most n, there is a pure Cohen-Macaulay S-module of type $\mathbf{d}$. This article is motivated by this conjecture, and the fact that it is true(see Remark 1.3(a)), as shown by D. Eisenbud and F. Schreyer.

In this paper, we study modules of finite projective dimension, with pure resolutions, over an arbitrary standard graded ring R. The main tool used is the Cohen-Macaulay defect of an R-module M, denoted $\operatorname{cmd}(M)$. We show that the existence of a pure R-module M of finite projective dimension forces $\operatorname{cmd}(R) \leq \operatorname{cmd}(M)$, a crucial observation that is used to prove the main results in this article. In particular, this shows that if such a module is Cohen-Macaulay, then the ring is also forced to be so.

In [1], Boij and Söderberg characterize graded Cohen-Macaulay modules over S, by relations among the graded Betti numbers of the module called the Herzog-Kühl equations. In Section 2, we generalize this result to arbitrary Cohen-Macaulay standard graded algebras in Theorem 2.6. The Boij-Soderberg result extends a theorem of J. Herzog and M. Kühl from [5], in which they prove it for pure Cohen-Macaulay modules over S. In Theorem 2.9, we generalize the result of Herzog and Kühl to a standard graded ring R, and characterize pure R-modules M satisfying $\operatorname{cmd}(M) = \operatorname{cmd}(R)$.

The above results are used in Section 3, first to compute the multiplicity of such modules in Corollary 3.1, and then to get some interesting observations about cyclic modules with pure resolutions in Subsection 3.2. In Subsection 3.3, we give a sufficient condition for a pure module over the polynomial ring to be Cohen-Macaulay. In particular, in Corollary 3.7, we see conditions on the degree sequences that force a pure quotient of a polynomial ring to be Gorenstein, or a complete intersection.

Finally, we see a characterization for R to be Cohen-Macaulay in terms of pure modules. Using Proposition 2.7, we prove (in Theorem 3.9) that R is Cohen-Macaulay if and only if there is a pure Cohen-Macaulay R-module corresponding to any degree sequence of length at most $\operatorname{depth}(R)$.

2010 *Mathematics Subject Classification.* Primary 13A02, 13C14, 13D02.

Key words and phrases. Herzog-Kühl equations, Cohen-Macaulay defect, graded Betti numbers, pure modules.

Section 1 contains the notation, definitions, basic observations, and previous results that are needed in the rest of the article.

1. Preliminaries

1.1. **Notation.**

a) By k, we mean an infinite field, and R denotes a standard graded k-algebra, i.e., $R_0 = \mathsf{k}$, and R is generated as an algebra over k by finitely many elements of degree 1.

b) M denotes a finitely generated graded R-module, $\mu(M)$, the *minimal number of generators* of M, $\operatorname{cmd}(M) = \dim(M) - \operatorname{depth}(M)$ is the *Cohen-Macaulay defect of* M, and the ith *total Betti number* of M is $\beta_i^R(M) = \dim_{\mathsf{k}}(\operatorname{Tor}_i^R(\mathsf{k}, M))$.

c) The (i,j)th-graded Betti number of M is $\beta_{i,j}^R(M) = \dim_{\mathsf{k}}(\operatorname{Tor}_i^R(M, \mathsf{k})_j)$. The *Betti diagram (or Betti table)* of M is written as

$\beta^R(M)=$

j \ i	0	1	$\cdots$	p	$\cdots$
$\vdots$	$\vdots$	$\vdots$	$\ddots$	$\vdots$	$\ddots$
0	$\beta_{0,0}$	$\beta_{1,1}$	$\cdots$	$\beta_{p,p}$	$\vdots$
1	$\beta_{0,1}$	$\beta_{1,2}$	$\cdots$	$\beta_{p,p+1}$	$\vdots$
$\vdots$	$\vdots$	$\vdots$	$\ddots$	$\vdots$	$\ddots$

, i.e., the (i,j)th entry is $\beta_{i,i+j}^R(M)$.

d) If M_j is the jth graded component of M, the *shifted module* $M(d)$ is the R-module M with graded components $M(d)_j = M_{j+d}$. The *Hilbert series* of M is $H_M(z) = \sum_{j\in\mathbb{Z}} \lambda(M_j) z^j$.

e) By the Hilbert-Serre Theorem (see [3, Section 4.1]), if $\dim(M) = r$, then there exists a unique Laurent polynomial $f(z)$ over $\mathbb{Z}$ with $f(1) > 0$ such that $H_M(z) = \frac{f(z)}{(1-z)^r}$. The *multiplicity* of M is $e(M) = f(1)$.

1.2. **Pure Modules and Degree Sequences.**

Definition 1.1. Let R be standard graded, and M a finitely generated graded R-module.

a) An R-module M has a *pure resolution* if each column of the Betti diagram of M has at most one non-zero entry. If this happens, we say that M is a *pure R-module*.

If M has a pure resolution, and $\beta_{i,d_i} \neq 0$ for each $i < \operatorname{pdim}_R(M) + 1$, then M is said to have a *pure resolution of type* $\mathbf{d} = (d_0, d_1, \ldots, d_p)$ if M is a module of finite projective dimension p over R, and of type $(d_0, d_1, \ldots)$ otherwise.

b) For $p \leq \operatorname{depth}(R)$, a sequence $\mathbf{d} = (d_0, d_1, \ldots, d_p)$ is said to be a *degree sequence* of length p if $d_i \in \mathbb{Z}$, and $d_i < d_{i+1}$ for all i.

In the following remark, we record the relation between the Hilbert series, Betti numbers and shifts in the minimal free resolution of a pure module.

Remark 1.2 (Hilbert series and Betti tables of pure modules)**.**
Let R be standard graded with Hilbert series $H_R(z) = \sum_{i\geq 0} h_i^R z^i$, M a pure R-module of type $\mathbf{d} = (d_0, d_1, \ldots)$, with Hilbert series $H_M(z) = \sum_{i\in\mathbb{Z}} h_i^M z^i$, and $\beta_i = \beta_{i,d_i}^R(M)$. Then the Hilbert series and Betti table of M can be obtained from one another as follows:

a) Fix $0 \leq i < \operatorname{pdim}_R(M)+1$. If K is the ith syzygy of M in a minimal R-free resolution, then the exact sequence

$$0 \to K \to R(-d_{i-1})^{\oplus \beta_{i-1}} \to \cdots \to R(-d_0)^{\oplus \beta_0} \to M \to 0$$

yields $H_K(z) = \left(\sum_{k=0}^{i-1}(-1)^{i-1-k}\beta_k z^{d_k}\right) H_R(z) + (-1)^i H_M(z)$.

Now, d_i is the order of $H_K(z)$, and β_i is the coefficient of z^{d_i} in $H_K(z)$. Hence, we see that d_i, and β_i can be inductively computed from $H_M(z)$ as

$$d_i = \min\left\{ j : \sum_{k=0}^{i-1}(-1)^k \beta_k h^R_{j-d_k} - h^M_j \neq 0 \right\}, \qquad \beta_i = \left| \sum_{k=0}^{i-1}(-1)^k \beta_k h^R_{d_i-d_k} - h^M_{d_i} \right|.$$

b) The additivity of Hilbert series on exact sequences applied to a minimal R-resolution of M shows that h^M_i can be obtained from d_i, and β_i by using $H_M(z) = H_R(z)\left(\sum_{i\geq 0}(-1)^i \beta_i z^{d_i}\right)$.

Remark 1.3 (Pure modules over polynomial rings)**.**
Let $S = \mathsf{k}[X_1, \ldots, X_n]$, and $\mathbf{d} = (d_0, d_1, \ldots, d_p)$, $p \leq n$, be a degree sequence.

a) (Eisenbud-Schreyer, [4, Theorem 5.1])
There exists a pure Cohen-Macaulay S-module of type $\mathbf{d}$.

b) (Herzog-Kühl, [5, Theorem 1])
If M is a pure S-module of type $\mathbf{d}$, then M is Cohen-Macaulay if and only if

$$\beta^S_i(M) = \beta^S_0(M) \prod_{j\neq 0,i} \left| \frac{d_j - d_0}{d_j - d_i} \right| \qquad \text{for } 1 \leq i \leq p.$$

c) We define two *pure diagrams* corresponding to $\mathbf{d}$:

$$\pi(\mathbf{d}) = \frac{1}{\beta^S_0(M)}\beta^S(M), \text{ and } \pi'(\mathbf{d}) = \frac{1}{\beta^S_p(M)}\beta^S(M),$$

where M is a pure Cohen-Macaulay S-module of type $\mathbf{d}$, which exists by (a).

By (b), these are independent of S and M, and for $0 \leq i \leq p$, and $j \in \mathbb{Z}$, we have

$$\pi(\mathbf{d})_{i,j} = \begin{cases} \prod_{k\neq i,0} \left|\frac{d_k-d_0}{d_k-d_i}\right| & \text{for } j = d_i \\ 0 & \text{for } j \neq d_i \end{cases} \quad \text{and} \quad \pi'(\mathbf{d})_{i,j} = \begin{cases} \prod_{k\neq i,p} \left|\frac{d_k-d_p}{d_k-d_i}\right| & \text{for } j = d_i \\ 0 & \text{for } j \neq d_i \end{cases}.$$

d) Boij-Söderberg([2, Theorem 2]) and Eisenbud-Schreyer([4, Theorem 7.1]) prove:

Let M be an S-module with $\operatorname{codim}(M) = c$ and $\operatorname{pdim}_S(M) = p$. There exist finitely many degree sequences $\mathbf{d}$ of length s, with $c \leq s \leq p$, such that $\beta^S(M) = \sum c_{\mathbf{d}}\pi(\mathbf{d})$, where $c_{\mathbf{d}} \in \mathbb{Q}_+$.

Since $\pi(\mathbf{d})$, and $\pi'(\mathbf{d})$ are positive rational multiples of each other for each $\mathbf{d}$, we see that there exist $c'_{\mathbf{d}} \in \mathbb{Q}_+$, such that $\beta^S(M) = \sum c'_{\mathbf{d}}\pi'(\mathbf{d})$.

2. Herzog-Kühl Equations

2.1. **Herzog-Kühl Equations over Arbitrary Rings.** The Herzog-Kühl equations are relations among the graded Betti numbers of a module over a polynomial ring. These were initially studied in [5], and then in [1], to characterize Cohen-Macaulay modules. We look at similar relations over all standard graded k-algebras in this section.

Definition 2.1. *Let R be a standard graded k-algebra, and M a graded R-module of finite projective dimension p. Let $\beta_{i,j} = \beta^R_{i,j}(M)$. We say that M satisfies the Herzog-Kühl equations if*

$$\sum_{i,j}(-1)^i j^l \beta_{i,j} = 0 \quad \text{for } l = 0, \ldots, p-1.$$

Remark 2.2. Let $S = \mathsf{k}[X_1, \ldots, X_n]$, and M be a graded S-module with $\operatorname{pdim}_S(M) = p$.
a) If M is pure of type $(d_0, \ldots, d_p)$, then M satisfies the Herzog-Kühl equations if and only if $\beta^S_i(M) = \beta^S_0(M) \prod_{j\neq 0,i} \left|\frac{d_j - d_0}{d_j - d_i}\right|$ for $1 \leq i \leq p$. (For an explanation, see [1], or the proof of the equivalence of (iii) and (iv) in Theorem 2.9). In light of Remark 1.3(b), the relations in Definition 2.1 are called the Herzog-Kühl equations.
b) In [1, Section 2.1], Boij and Söderberg extend the Herzog-Kühl result to S-modules that are not necessarily pure. They prove that an S-module M satisfies the Herzog-Kühl equations in Definition 2.1 if and only if it is Cohen-Macaulay.

We begin with the following observation:

Remark 2.3. Let $h(z) = \sum_{i,j}(-1)^i b_{i,j} z^j$ be a Laurent polynomial over $\mathbb{Z}$. Then $(1-z)^n$ divides $h(z)$ if and only if the $b_{i,j}$'s satisfy the following equations:

$$\sum_{i,j}(-1)^i j^l b_{i,j} = 0 \quad \text{for } l = 0, \ldots, n-1.$$

In order to see this, let $a_l = \sum_{i,j}(-1)^i j^l b_{i,j} = 0$ for $l = 0, \ldots, n-1$. Then $h(1) = a_0$, and

$$h^{(l)}(1) = a_l - \left(\sum_{1\leq k\leq l} k\right) a_{l-1} + \left(\sum_{1\leq k_1<k_2\leq l} k_1 k_2\right) a_{l-2} - \cdots + (-1)^{l-1}\left(\prod_{k=1}^{l} k\right) a_1 \text{ for } 1 \leq l < n.$$

Thus $\begin{pmatrix} h(1) \\ h^{(1)}(1) \\ \vdots \\ h^{(n-1)}(1) \end{pmatrix} = A \begin{pmatrix} a_0 \\ a_1 \\ \vdots \\ a_{n-1} \end{pmatrix}$, where A is an $n \times n$ lower triangular matrix with diagonal entries 1.

Thus we are done, since $(1-z)^n$ divides $h(z)$ if and only if $h^{(l)}(1) = 0$ for $0 \leq l < n$, which, by the invertibility of A, happens if and only if $a_0, \ldots, a_{n-1}$ are all 0.

Proposition 2.4. *Let R be standard graded, and M a graded R-module of codimension c and finite projective dimension p. If $\beta_{i,j} = \beta^R_{i,j}(M)$, and $h(z) = \sum_{i=0}^{p}\sum_{j\in\mathbb{Z}}(-1)^i \beta_{i,j} z^j$, then:*

a) *$(1-z)^c$ divides $h(z)$, and $(1-z)^{c+1}$ does not divide $h(z)$,*
b) *the graded Betti numbers of M satisfy:*

$$\sum_{i=0}^{p}\sum_{j\in\mathbb{Z}}(-1)^i j^l \beta_{i,j} = 0 \quad \text{for } l = 0, \ldots, c-1, \text{ and}$$

c) *the multiplicity of M is $e(M) = \dfrac{e(R)}{c!}\left[\sum_{i=0}^{p}\sum_{j\in\mathbb{Z}}(-1)^{i+c} j^c \beta_{i,j}\right]$.*

Proof. The proof of (b) follows from (a) and the previous remark. Let $\dim(R) = d$. Hence, the Hilbert series of R and M are of the form $H_R(z) = \frac{f(z)}{(1-z)^d}$, and $H_M(z) = \frac{g(z)}{(1-z)^{d-c}}$, where $e(R) = f(1) \neq 0 \neq g(1) = e(M)$.

Since a minimal resolution of M over R is of the form

$$0 \to \bigoplus_j R(-j)^{\beta_{p,j}} \to \cdots \to \bigoplus_j R(-j)^{\beta_{1,j}} \to \bigoplus_j R(-j)^{\beta_{0,j}} \to M \to 0,$$

by the additivity of Hilbert series over exact sequences, we get

$$\frac{g(z)}{(1-z)^{d-c}} = H_M(z) = H_R(z) \left(\sum_{i=0}^{p} \sum_{j \in \mathbb{Z}} (-1)^j \beta_{i,j} z^j \right) = \frac{f(z)h(z)}{(1-z)^d}.$$

Thus, $g(z)(1-z)^c = f(z)h(z)$. Since $f(1) \neq 0$, we see that $(1-z)^c$ divides $h(z)$, and $g(1) \neq 0$ implies that $(1-z)^{c+1}$ does not divide $h(z)$, proving (a), and hence (b).

In order to prove (c), let $h(z) = (1-z)^c h_1(z)$. Therefore, $h_1(1) =$

$$\frac{(-1)^c}{c!} h^{(c)}(1) = \frac{(-1)^c}{c!} \left(\sum_{i,j} (-1)^i j(j-1)(j-2)\cdots(j-c+1)\beta_{i,j} \right) = \frac{1}{c!} \sum_{i,j} (-1)^{i+c} j^c \beta_{i,j},$$

where the last equality is by (b). Thus, $e(M) = g(1) = e(R)h_1(1)$, proving (c). □

Remark 2.5. Let R be a standard graded k-algebra, and M be an R-module of finite projective dimension p and $\operatorname{codim}(M) = c$. By Remark 2.3 and Proposition 2.4, we see that M satisfies the Herzog-Kühl equations if and only if $p \leq c$. By the Auslander-Buchsbaum formula, since $\operatorname{pdim}_R(M) < \infty$, this is equivalent to $\operatorname{cmd}(M) \leq \operatorname{cmd}(R)$.

Observe that if R is Cohen-Macaulay in the above remark, i.e., $\operatorname{cmd}(R) = 0$, then M satisfies the Herzog-Kühl equations if and only if $\operatorname{cmd}(M) = 0$, i.e., M is Cohen-Macaulay. Thus we have proved:

Theorem 2.6. *Let R be a standard graded Cohen-Macaulay k-algebra and M an R-module of finite projective dimension p. Then M is Cohen-Macaulay if and only if its graded Betti numbers satisfy the equations*

$$\sum_{i=0}^{p} \sum_{j \in \mathbb{Z}} (-1)^i j^l \beta_{i,j} = 0, \quad l = 0, \ldots, p-1.$$

In particular, the above theorem extends the Boij-Söderberg result in Remark 2.2(b), characterizing graded Cohen-Macaulay R-modules by the Herzog-Kühl equations.

2.2. **Herzog-Kühl Equations for Pure Modules.** We now show, in Theorem 2.9, that the Herzog-Kühl equations can be used to characterize pure modules M of finite projective dimension over a standard graded k-algebra R, satisfying $\operatorname{cmd}(M) = \operatorname{cmd}(R)$. This generalizes the theorem of Herzog and Kühl in Remark 1.3(b), and extends Theorem 2.6 to pure modules over a standard graded k-algebra.

We now prove a key proposition, which is used in this section and the next.

Proposition 2.7. *Let R be a standard graded k-algebra, and M a pure R-module. Then* $\operatorname{codim}(M) \leq \operatorname{pdim}_R(M)$. *Moreover, if* $\operatorname{pdim}_R(M)$ *is finite, then* $\operatorname{cmd}(R) \leq \operatorname{cmd}(M)$.

Proof. Note that if $\operatorname{pdim}_R(M)$ is infinite, there is nothing to prove. Hence, assume that M is pure of type $\mathbf{d} = (d_0, \ldots, d_p)$, in particular, assume that $\operatorname{pdim}_R(M) = p$ is finite. With $\operatorname{codim}(M) = c$, we want to prove $c \leq p$.

If $p < c$, then, by Proposition 2.4(b), $y_i = \beta_{i,d_i}$, $i = 0, \ldots, p$ is a non-trivial solution of the linear system of equations:

$$\sum_i (-1)^i d_i^l y_i = 0, \qquad l = 0, 1, \ldots, c-1. \tag{1}$$

The maximal minors of the coefficient matrix A of this system are $(p+1) \times (p+1)$ Vandermonde matrices since $p + 1 \leq c$. Thus, $d_i \neq d_j$ for all $i \neq j$ implies that A has maximum rank $p + 1$. This forces the system of equations (1) to have only the trivial solution. This contradiction shows that $c \leq p$.

Finally, since $\operatorname{pdim}_R(M)$ is finite, then the Auslander-Buchsbaum formula implies that $\operatorname{codim}(M) \leq \operatorname{pdim}_R(M)$ is equivalent to $\operatorname{cmd}(R) \leq \operatorname{cmd}(M)$, proving the result. □

In the following example, we see that the inequality $\operatorname{cmd}(R) \leq \operatorname{cmd}(M)$ can be strict.

Example 2.8. Let $S = \mathsf{k}[X_1, X_2]$, and $M = S/\langle X_1^2, X_1X_2\rangle$. Then M is pure of type $\mathbf{d} = (0, 2, 3)$. In this case, $\operatorname{cmd}(M) = 1 > 0 = \operatorname{cmd}(S)$.

Note that $\beta^R(M)$ does not satisfy the Herzog-Kühl equations, since

$$\beta^R(M) = \begin{array}{|c|ccc|}\hline j \backslash i & 0 & 1 & 2 \\ \hline 0 & 1 & - & - \\ 1 & - & 2 & 1 \\ \hline \end{array} \quad \neq \quad \begin{array}{|c|ccc|}\hline j \backslash i & 0 & 1 & 2 \\ \hline 0 & 1 & - & - \\ 1 & - & 3 & 2 \\ \hline \end{array} = \beta_0^S(M)\pi(\mathbf{d}).$$

Theorem 2.9. *Let M be a pure module of type $\mathbf{d} = (d_0, \ldots, d_p)$ over a standard graded k-algebra R. Let $\pi(\mathbf{d})$ and $\pi'(\mathbf{d})$ be as in Remark 1.3(c). Then the following are equivalent:*
i) $\operatorname{cmd}(M) = \operatorname{cmd}(R)$.
ii) $\operatorname{codim}(M) = \operatorname{pdim}_R(M)$.
iii) $\sum_{i=0}^{p} (-1)^i d_i^l \beta_{i,d_i} = 0$ *for* $l = 0, \ldots, p-1$, *i.e. M satisfies the Herzog-Kühl equations.*
iv) $\beta^R(M) = \beta_0^R(M)\pi(\mathbf{d})$.
v) $\beta^R(M) = \beta_p^R(M)\pi'(\mathbf{d})$.

Proof. Since M is a pure module of type $\mathbf{d}$, we have $\operatorname{pdim}_R(M) = p$ is finite. Hence, by the Auslander-Buchsbaum formula, (i) and (ii) are restatements of each other.

Let $\operatorname{codim}(M) = c$. If (ii) is true, then Proposition 2.4(b) shows that (iii) holds. In order to prove the converse, first note that by Remark 2.5, (iii) implies that $p \leq c$. Furthermore, since M is a pure R-module, by Proposition 2.7, we get $c \leq p$, proving (ii).

In order to see the equivalence of (iii) and (iv), note that the Herzog-Kühl equations are p linear equations in $\beta_{0,d_0}, \ldots, \beta_{p,d_p}$, whose coefficient matrix is $\begin{bmatrix} 1 & \cdots & (-1)^p \\ d_0 & \cdots & (-1)^p d_p \\ \vdots & \ddots & \vdots \\ d_0^{p-1} & \cdots & (-1)^p d_p^{p-1} \end{bmatrix}$. Since $d_i \neq d_j$ for $i \neq j$, this a $p \times (p+1)$ matrix of full rank, and the solutions can be obtained by setting β_{0,d_0} to be a free variable. Cramer's Rule proves the rest.

Similarly, setting β_{p,d_p} to be a free variable proves the equivalence of (iii) and (v). □

3. Applications

3.1. **Application to Multiplicity.** In [1], Boij and Söderberg compute the multiplicity of pure Cohen-Macaulay modules over polynomial rings. We use the above theorem to do the same for a pure module M satisfying $\operatorname{pdim}_R(M) = \operatorname{codim}(M)$, over any standard graded k-algebra R.

Corollary 3.1. *Let M be a pure module of type $\mathbf{d} = (d_0, \ldots, d_p)$ over a standard graded k-algebra R, with $\beta_0 = \beta^R_{0,d_0}(M)$. If $\operatorname{cmd}(R) = \operatorname{cmd}(M)$, then the multiplicity of M is*

$$e(M) = e(R)\frac{\beta_0}{p!}\prod_{j=1}^{p}(d_j - d_0).$$

Proof. Since $\operatorname{cmd}(R) = \operatorname{cmd}(M)$, and $\operatorname{pdim}_R(M) = p$ is finite, we get $\operatorname{codim}(M) = p$. Hence, by Theorem 2.9 and Proposition 2.4, we have

$$\begin{aligned} e(M) &= \beta_0\frac{e(R)}{p!}\sum_{i=0}^{p}(-1)^{i+p}d_i^p\prod_{j\neq i, j=1}^{p}\left|\frac{d_j - d_0}{d_j - d_i}\right| \\ &= e(R)\frac{\beta_0}{p!}\prod_{j=1}^{p}(d_j - d_0)\left(\sum_{i=0}^{p}\frac{(-1)^{i+p}d_i^p}{\prod\limits_{j\neq i, j=0}^{p}|d_j - d_i|}\right). \end{aligned}$$

Let V be the $(p+1)\times(p+1)$ Vandermonde matrix with (i,j)th entry d_{i-1}^{j-1}. We know that $\det(V) = \prod\limits_{0\leq j<k\leq p}(d_k - d_j)$. On the other hand, expanding along the last row, we get

$$\det(V) = \sum_{i=0}^{p}(-1)^{i+p}d_i^p\left(\prod_{0\leq j<k\leq p,\ j\neq i\neq k}(d_k - d_j)\right).$$

Dividing the second expression for $\det(V)$ by the first, we see that $\sum\limits_{i=0}^{p}(-1)^{i+p}\frac{d_i^p}{\prod\limits_{j\neq i, j=0}^{p}|d_j - d_i|} = 1$. Substituting this in the expression for $e(M)$ proves the corollary. □

3.2. **Applications to Cyclic Modules.** In this subsection, we study pure cyclic modules. We first look at Betti diagrams of pure cyclic Cohen-Macaulay modules of projective dimension 2.

Theorem 3.2. *Let R be a standard graded k-algebra, and I a homogeneous ideal in R, such that $\operatorname{codim}(R/I) = \operatorname{pdim}_R(R/I) = 2$. If R/I is a pure R-module, then there exists a regular sequence g_1, g_2 such that $\beta^R(R/I) = \beta^R(R/\langle g_1, g_2\rangle^i)$ for some i.*

Proof. Let R/I be a pure R-module of type $(0, d_1, d_2)$. Since $\operatorname{codim}(R/I) = \operatorname{pdim}_R(R/I)$ and $\beta_0^R(R/I) = 1$, by Theoqrem 2.9, we have $\beta_1^R(R/I) = d_2/(d_2 - d_1)$ and $\beta_2^R(R/I) = d_1/(d_2 - d_1)$. Therefore $r = d_2 - d_1 = \gcd(d_1, d_2)$.

Write $d_1 = rd$ and let g_1, g_2 be an R-regular sequence of degree r. By using induction on d, it can be seen that $R/\langle g_1, g_2\rangle^d$ is a pure R-module of type $(0, rd, rd + r)$. Therefore $R/\langle g_1, g_2\rangle^d$ and R/I are pure of type $(0, rd, rd + r)$, with $\beta_0^R(R/I) = \beta_0^R(R/\langle g_1, g_2\rangle^d) = 1$. Hence, by Theorem 2.9, $\beta^R(R/\langle g_1, g_2\rangle^d) = \beta^R(R/I)$. □

In the above theorem, we have seen that $\beta^R(R/\langle g_1, g_2\rangle^d) = \beta^R(R/I)$. One can ask: Is $R/I \simeq R/\langle g_1, g_2\rangle^d$? The following example shows that it need not be true.

Example 3.3. Let $S = \mathsf{k}[X_1, X_2, X_3]$, and $I = \langle X_1X_2, X_2X_3, X_1X_3\rangle$. Then S/I is a pure S-module of type $(0, 2, 3)$, and $\beta^S(S/I) = \beta^S(S/\langle X_1, X_2\rangle^2)$, illustrating Theorem 3.2.

We know that if g_1, g_2 is a regular sequence, then $S/\langle g_1, g_2\rangle^i$ is a Cohen-Macaulay for all $i \in \mathbb{N}$. Since S/I^2 is not Cohen-Macaulay, there does not exist any regular sequence g_1, g_2 such that I isomorphic to $\langle g_1, g_2\rangle^i$.

The following example shows that Theorem 3.2 need not hold if $\operatorname{pdim}_R(R/I) \geq 3$.

Example 3.4. Let $S = \mathsf{k}[X_1, X_2, X_3]$ and $I = \langle X_1X_2, X_2X_3, X_1X_3, X_1^2 - X_2^2, X_1^2 - X_3^2\rangle$. Then S/I is a pure Gorenstein quotient of S of type $(0, 2, 3, 5)$.

Note that $\beta^S(S/I) \neq \beta^S(S/\langle g_1, g_2, g_3\rangle^i)$, for any i, where g_1, g_2, g_3 is an S-regular sequence. This follows since $\beta_1^S(S/I) = 5$, and for any regular sequence g_1, g_2, g_3, we have $\beta_1^S(S/\langle g_1, g_2, g_3\rangle) = 3$, and $\beta_1^S(S/\langle g_1, g_2, g_3\rangle^i) \geq 6$ for $i \geq 2$.

We have seen in Theorem 3.2 that $\beta^R(R/I) = \beta^R(R/\langle g_1, g_2\rangle^i)$. Since $R/\langle g_1, g_2\rangle^i$ is a pure module for each i, we have the following natural question.

Question 3.5. Let R be a graded k-algebra and I be a homogeneous ideal in R such that R/I is a pure R-module. If $\operatorname{cmd}(R/I) = \operatorname{cmd}(R)$, then is R/I^i a pure R-module for every i?

3.3. **Modules over Polynomial Rings.** We first give a sufficient condition for a pure module over a polynomial to be Cohen-Macaulay. This leads to conditions on the degree sequences, which force pure cyclic quotients to be Gorenstein, or a complete intersection.

Theorem 3.6. *Let S be a polynomial ring, $\mathbf{d} = (d_0, d_1, \ldots, d_p)$ be a degree sequence with $1 \leq \pi'(\mathbf{d})_{0,d_0}$, and M a pure S module of type $\mathbf{d}$. If $\beta_{0,d_0}(M) \leq \beta_{p,d_p}(M)$, then M is Cohen-Macaulay.*

Proof. By Remark 1.3(d), we can write

$$\beta^S(M) = c_1\pi'(\mathbf{d}) + c_2\pi'(d_0, d_1, \ldots, d_{p-1}) + \cdots + c_{p-1}\pi'(d_0, d_1) + c_p\pi'(d_0),$$

where $c_i \in \mathbb{Q}_{\geq 0}$, with $c_1 > 0$. The above equation shows that $c_1 = \beta^S_{p,d_p}(M)$. Now

$$\begin{aligned}\beta^S_{0,d_0}(R) &= c_1\pi'(\mathbf{d})_{0,d_0} + c_2\pi'(d_0, d_1, \ldots, d_{p-1})_{0,d_0} + \cdots + c_{p-1}\pi'(d_0, d_1)_{0,d_0} + c_p\pi'(d_0)_{0,d_0}\\ &\geq c_1\pi'(\mathbf{d})_{0,d_0} \geq c_1 = \beta^S_{p,d_p}(M) \geq \beta^S_{0,d_0}(M).\end{aligned}$$

Thus $\pi'(\mathbf{d})_{0,d_0} = 1$, $\beta^S_{0,d_0}(M) = \beta^S_{p,d_p}(M)$. Moreover, $\beta^S(M) = \beta^S_{p,d_p}(M)\pi'(\mathbf{d})$, since $c_i = 0$ for $i \geq 2$. Therefore, M is Cohen-Macaulay by Remark 1.3(b) and (c). □

Corollary 3.7. *Let S be a polynomial ring, $\mathbf{d} = (0, d_1, \ldots, d_p)$ be a degree sequence, and $\mathbf{e} = (e_1, \ldots, e_p)$, where $e_i = d_i - d_{i-1}$ for $i = 1, \ldots, p$. Let $I \subset S$ be homogeneous such that $R = S/I$ is a pure S-module of type $\mathbf{d}$. Then the following hold:*

a) *If $\pi'(\mathbf{d})_{0,d_0} \geq 1$, then R is Gorenstein.*

b) *If $\mathbf{e}$ is non-decreasing, then $e_1 = \cdots = e_p$, and R is a complete intersection.*

Proof. a) Note that R is Gorenstein if and only if R is Cohen-Macaulay, and $\beta^S_p(R) = 1$. Both the latter conditions hold by the previous theorem, hence R is Gorenstein.

b) We have $d_p = \sum_{i=1}^p e_i = \sum_{i=1}^j e_i + \sum_{i=j+1}^p e_i \geq \sum_{i=1}^j e_i + \sum_{i=1}^{p-j} e_i = d_j + d_{p-j}$, for each j, since $\mathbf{e}$ is non-decreasing. In particular, $\frac{d_p - d_j}{d_{p-j}} \geq 1$, with equality holding for each j if and only if $e_1 = \cdots = e_p$.

Now, by definition, $\pi'(\mathbf{d})_{0,0} = \prod_{j=1}^{p-1}\left(\frac{d_p - d_j}{d_j}\right) = \prod_{j=1}^{p-1}\left(\frac{d_p - d_j}{d_{p-j}}\right) \geq 1$. Hence, by the previous theorem, $\pi'(\mathbf{d})_{0,0} = 1$, which forces $\frac{d_p - d_j}{d_{p-j}} = 1$ for each j, and hence, we see that $e_1 = \cdots = e_p$.

Thus, if $d_1 = d$, then $\mathbf{d} = (0, d, 2d, \ldots, pd)$. We see that I is generated by a regular sequence, since $\mu(I) = \beta_1^S(R) = \beta_0^S(R)\prod_{k=2}^{p}\left|\frac{d_k}{d_k - d_1}\right| = p = \operatorname{codim}(R)$. Thus R is a complete intersection. □

3.4. **Characterization of Cohen-Macaulay Rings.** In Theorem 3.9, we give a characterization of Cohen-Macaulay graded k-algebras in terms of pure Cohen-Macaulay modules. We first prove the following proposition.

Proposition 3.8. *Let R be a standard graded Cohen-Macaulay k-algebra of dimension n, and $\mathbf{d} = (d_0, d_1, \ldots, d_p)$, $p \leq n$, be a degree sequence. Then there exists a pure Cohen-Macaulay R-module of type $\mathbf{d}$.*

Proof. Let $\operatorname{depth}(R) = r$. Since k is infinite, there exists a regular sequence $l_1, \ldots, l_r$ of linear forms in R. Since R is Cohen-Macaulay, R is free of finite rank over $S = \mathsf{k}[l_1, \ldots, l_r]$, which is isomorphic to a polynomial ring over k.

By Remark 1.3(a), there exists a Cohen-Macaulay S-module M, which has a pure resolution over S of type $\mathbf{d}$. Then $M \otimes_S R$ is a pure Cohen-Macaulay R-module of type $\mathbf{d}$. □

The next theorem follows from Propositions 2.7 and 3.8.

Theorem 3.9. *Let R be a standard graded k-algebra. Then the following are equivalent:*
i) *there exists a pure Cohen-Macaulay R-module of type $\mathbf{d}$, for every degree sequence $\mathbf{d}$ of length at most* $\operatorname{depth}(R)$.
ii) *there exists a pure Cohen-Macaulay R-module of finite projective dimension.*
iii) *R is Cohen-Macaulay.*

Proof. Statement (ii) follows immediately from (i). Since M is Cohen-Macaulay if and only if $\operatorname{cmd}(M) = 0$, the proof of (ii) ⇒ (iii) is a consequence of Proposition 2.7, and (iii) ⇒ (i) is the content of Proposition 3.8. □

References

[1] M. Boij and J. Söderberg, *Graded Betti Numbers of Cohen-Macaulay Modules and the Multiplicity Conjecture,* J. Lond. Math. Soc.(2) **78** (2008), no. 1, 85–106.

[2] M. Boij and J. Söderberg, *Betti numbers of graded modules and the multiplicity conjecture in the non-Cohen-Macaulay case,* Algebra Number Theory **6** (2012), no. 3, 437–454.

[3] W. Bruns and J. Herzog, Cohen-Macaulay Rings. Cambridge Studies in Advanced Mathematics, **39**. Cambridge University Press, Cambridge, 1999.

[4] D. Eisenbud and F. Schreyer, *Betti Numbers of Graded Modules and Cohomology of Vector Bundles,* J. Amer. Math. Soc. **22**(2009), no. 3, 859–888.

[5] J. Herzog, and M. Kühl, *On the Betti Numbers of Finite Pure and Linear Resolutions,* Comm. Algebra **12** (1984), 1627–1646.

Department of Mathematics, I.I.T. Bombay, Powai, Mumbai 400076.
E-mail address: ananth@math.iitb.ac.in

Department of Mathematics, I.I.T. Bombay, Powai, Mumbai 400076.
E-mail address: rajiv@math.iitb.ac.in